

An important issue in string theory is the study of threshold corrections in gauge couplings [1]. In particular their dependence on the moduli of the compactified space plays a crucial role in string unification, the derivation of the low energy effective Lagrangian, and the problem of supersymmetry breaking. This dependence was studied extensively in the case of symmetric orbifold compactification of the heterotic superstring for untwisted moduli [1],[2], and there was found that it satisfies a non renormalization theorem [3]. Namely, it is given entirely at one-loop level and it is determined by the violation of the integrability condition of the corresponding Θ -angles with respect to moduli. This phenomenon has also been understood in the field theory framework as an infrared effect due to propagation of massless fermions in one-loop anomalous graphs [2],[4]. A general derivation, from the point of view of the internal $N = 2$ superconformal theory, is presented in [5].

In this note we extend the analysis of [1],[2],[3], to the study of threshold corrections in gravitational couplings. In analogy to the case of gauge couplings, to avoid infrared divergences, we consider the on-shell three point function of two gravitons $h_{\mu\nu}$, $h_{\lambda\rho}$ and one modulus field T , with momenta p_1, p_2 and p_3 respectively:

$$\langle h^{\mu\nu}(p_1)h^{\lambda\rho}(p_2)T(p_3) \rangle = \int \langle V_h^{\mu\nu}(p_1, z_1)V_h^{\lambda\rho}(p_2, z_2)V_T(p_3, z_3) \rangle. \quad (1)$$

In (1) the integration is extended over the moduli space of the world-sheet surface as well as over the positions of the graviton and modulus vertex operators V_h and V_T . There is also an implicit summation over all even or odd spin structures corresponding to the CP-even or CP-odd part of the amplitude. The vertex operators for genus $g \geq 1$ read:

$$V_h^{\mu\nu} =: \bar{\partial}X^\mu(\partial X^\nu + ip \cdot \psi\psi^\nu)e^{ip \cdot X} :, \quad (2a)$$

$$V_T = v_{IJ} : \bar{\partial}X^I(\partial X^J + ip \cdot \psi\psi^J)e^{ip \cdot X} :, \text{ CP even}, \quad (2b)$$

$$V_T = v_{IJ} : \bar{\partial}X^I\psi^J e^{ip \cdot X} :, \text{ } g = 1, \text{ CP odd}, \quad (2c)$$

$$v_{IJ} = \partial_{\langle T \rangle}(G_{IJ} + B_{IJ}), \quad (2d)$$

The ψ 's are the left moving fermionic superpartners of the bosonic coordinates X 's, and G_{IJ} and B_{IJ} are the metric and the two-index antisymmetric tensor of the six-dimensional internal space, respectively. The different form of the vertex operator V_T between (2b) and (2c) is due to the existence of a conformal Killing spinor for the genus one case of odd spin structures, which implies also a "picture changing" of the form (2b), as well as the inclusion of one supercurrent insertion T_F associated to the corresponding supermodulus:

$$T_F =: \psi_\mu \partial X^\mu + G_{KL} \psi^L \partial X^K : + \text{ghosts}. \quad (3)$$

Let us first observe that the two derivative part of the amplitude (1) vanishes for $g = 1$, because every graviton leg contributes at least two powers of external momenta. Consider for instance the zeroeth and first order terms in the momentum expansion of the graviton vertex operator $V_h^{\mu\nu}$

$$V_h^{\mu\nu} = \bar{\partial}_{z_1} X^\mu [\partial_{z_1} X^\nu + ip_1 \cdot X \partial_{z_1} X^\nu + ip_1 \cdot \psi \psi^\nu] + O(p_1^2). \quad (4)$$

Then, only the part of the amplitude where the ψ 's are contracted can in principle contribute, due to the summation over spin structures, and one is left essentially with a term of the kind:

$$\langle \bar{\partial}_{z_1} X(z_1) \bar{\partial}_{z_2} X(z_2) \rangle \langle (\psi(z_1) \psi(z_2)) \rangle^2.$$

Here again summation over spin structures removes the pole of the fermionic correlator and one can safely partial integrate, so that even this term actually gives vanishing contribution to (1).

Therefore, there is no T-dependent, one-loop correction to the Planck mass and the amplitude (1) contributes only to four derivative gravitational terms. These are quadratic in the Riemann tensor and their structure is very similar to that of the gauge kinetic terms¹:

¹ Unitarity arguments [6] forbid the appearance of other combinations of R_{abcd}^2, R_{ab}^2 , and R^2 .

$$L_{gr} = \Delta^{gr}(T, \bar{T}) R \wedge R^* + \Theta^{gr}(T, \bar{T}) R \wedge R. \quad (5)$$

The CP-even part $R \wedge R^* = R_{abcd}^2 - 4R_{ab}^2 + R^2$ is the topological Gauss-Bonnet combination whose integral gives the Euler number, the CP-odd part $R \wedge R = \epsilon^{abcd} R_{abef} R_{cd}^{ef}$ is the gravitational analog of $F \wedge F$. The on-shell three-point function (1) gives rise to $\Delta_T^{gr}(\langle T \rangle, \langle \bar{T} \rangle) \equiv \partial_{\langle T \rangle} \Delta^{gr}(\langle T \rangle, \langle \bar{T} \rangle)$ and Θ_T^{gr} , which are related by $N = 1$ supersymmetry as follows:

$$\partial_{\langle T \rangle} \Delta^{gr} + i\Theta_T^{gr} = 0, \quad \partial_{\langle \bar{T} \rangle} \Delta^{gr} - i\Theta_{\bar{T}}^{gr} = 0. \quad (6)$$

In fact, supersymmetry relates (1) to the three-point fermionic amplitude of a graviton, gravitino and fermion partner of T , whose chirality implies the complexification (6) of the original bosonic amplitude (1). Furthermore if $\Theta^{gr}(\langle T \rangle, \langle \bar{T} \rangle)$ is integrable, i.e. it satisfies:

$$\partial_{\langle \bar{T} \rangle} \Theta_T^{gr} - \partial_{\langle T \rangle} \Theta_{\bar{T}}^{gr} = 0, \quad (7)$$

then (6) implies that Δ^{gr} and Θ^{gr} are the real and imaginary parts respectively of an analytic function of the moduli T .

The similarity of the above expressions to those of the gauge kinetic terms allows one to extend the proof of the non renormalization theorem of [3] to the gravitational case. This can be done in a straightforward way in two steps: firstly all analytic corrections must vanish because they are inconsistent with duality invariance. Thus, the only possible corrections are those which violate the integrability condition (7). Secondly, the integrability condition is satisfied in all loops $g \geq 1$ pointwise in the world-sheet moduli, which shows that there are no higher loop corrections. At the one-loop level the calculation can be carried out explicitly or, alternatively, the result can be determined through the violation of the integrability condition, using again duality invariance. Below we present the one-loop calculation for the gravitational couplings in (5).

Following the same arguments used above to prove that each graviton contributes at least two powers of momenta, and using the on-shell relations $p_i \cdot p_j = 0$, it is easy to show that non vanishing contributions arise only if $\bar{\partial}_{z_1} X^\mu$ is contracted with $e^{ip_2 \cdot X(z_2)}$ and $\bar{\partial}_{z_2} X^\lambda$ with $e^{ip_1 \cdot X(z_1)}$, yielding the correlators:

$$\langle \bar{\partial}_{z_1} X^\mu(z_1) p_2 \cdot X(z_2) \rangle \langle \bar{\partial}_{z_2} X^\lambda(z_2) p_1 \cdot X(z_1) \rangle = -p_2^\mu p_1^\lambda \langle \bar{\partial}_{z_1} X(z_1) X(z_2) \rangle^2. \quad (8)$$

Also the pure bosonic part of the graviton vertex operator vanishes because of the summation over fermionic spin structures, yielding the fermionic correlators:

$$\langle p_1 \cdot \psi(z_1) \psi^\rho(z_2) \rangle \langle \psi^\nu(z_1) p_2 \cdot \psi(z_2) \rangle = \begin{cases} p_1^\rho p_2^\nu \langle \psi(z_1) \psi(z_2) \rangle^2, & CP \text{ even}, \\ -\epsilon^{\nu\rho\alpha\beta} p_{1,\alpha} p_{2,\beta}, & CP \text{ odd}. \end{cases} \quad (9)$$

Finally $\bar{\partial} X^I$ in V_T cannot be contracted and thus only its zero mode part contributes; the result is then proportional to:

$$v_{IJ} \bar{\partial} X^I \partial X^J, \quad CP \text{ even}, \quad (10a)$$

$$v_{IJ} G_{KL} \langle \psi^J \psi^L \rangle \bar{\partial} X^I \partial X^K, \quad CP \text{ odd}, \quad (10b)$$

where in (10b) the supercurrent insertion (3) was used.

From eqs. (10) it follows that non vanishing contribution is possible only from sectors which preserve $N = 2$ spacetime supersymmetry. Indeed, in $N = 1$ sectors all internal coordinates X^I are twisted by the action of the orbifold group and consequently they do not have zero modes. On the other hand, $N = 4$ sectors give vanishing contribution in the CP even part because of the fermionic spin-structure summation, and in the CP odd case because of the six internal fermionic zero-modes which cannot be saturated. In the $N = 2$ sectors, only one internal complex plane remains untwisted and X^I have momenta in the direction of the corresponding untwisted torus. Taking apart the momentum dependence and performing the two trivial z -integrations, one finds:

$$\Theta_T^{gr} = \frac{v_{IJ}\epsilon^{JL}G_{KL}}{4(2\pi)^4|G|^{1/2}} \int \frac{d^2z d^2\tau}{\tau_2^2} \langle \bar{\partial}X(z)X(0) \rangle^2 Z_{torus}^{IK} Z_{rest}(\bar{\tau}), \quad (11)$$

where $\tau = \tau_1 + i\tau_2$ is the modulus of the worldsheet torus, and

$$Z_{torus}^{IJ} = \sum_{p_L, p_R} p_L^I p_R^K e^{i\pi\tau p_L^2} e^{-i\pi\bar{\tau} p_R^2}$$

in terms of left and right moving momenta (p_L, p_R) of the untwisted complex plane, and we used the relation $\langle \psi^J \psi^L \rangle = \epsilon^{JL}$. Z_{rest} is purely antiholomorphic as a consequence of the cancellation between the contribution of the two trasverse left-moving spacetime coordinates X_L^μ and the two internal untwisted coordinates X_L^I , and the contribution of their fermionic superpartners ψ^ν and ψ^I . The last integration in (11) can be performed explicitly, giving the result:

$$\int d^2z \langle \bar{\partial}X(\bar{z})X(0) \rangle^2 = -2\pi i \tau_2 \frac{d}{d\tau_2} \log(\tau_2 \bar{\eta}^2(\tau)), \quad (12)$$

where $\eta(\tau) = q^{1/12} \prod_{n \geq 1} (1 - q^{2n})$, with $q = e^{i\pi\tau}$, is the Dedekind η -function. Substituting (12) into (11) and using the explicit form of the two dimensional lattice momenta, one finally finds:

$$\Theta_T^{gr} = \frac{i}{16\pi^2} \partial_{\langle T \rangle} \int \frac{d^2\tau}{\tau_2} Z_{torus} \frac{d}{2\pi i d\bar{\tau}} \log(\tau_2 \bar{\eta}^2) Z_{rest}(\bar{\tau}), \quad (13)$$

where $Z_{torus} = \sum_{p_L, p_R} q^{p_L^2} \bar{q}^{p_R^2}$.

Following the arguments of ref.[1], if one considers differences between the gravitational and gauge couplings, the τ_2 dependence inside the logarithm of (13) drops. Furthermore, the remaining antiholomorphic function $\frac{d}{2\pi i d\bar{\tau}} (\log \bar{\eta}^2) Z_{rest}$ is modular invariant and can have at most a singularity of the kind $\bar{q}^{-2}$ as $\tau_2 \rightarrow \infty$; if this is the case it implies that it is the j -invariant up to an additive constant, but in any case τ_1 integration picks the constant term, $\hat{b}^{gr}$, in its q -expansion. Δ^{gr} can then be determined using target space duality invariance from the violation of the integrability condition:

$$\partial_{\bar{T}} \Theta_T^{gr} - \partial_T \Theta_{\bar{T}}^{gr} = \frac{i \hat{b}^{gr}}{16\pi^2} \int \frac{d^2\tau}{\tau_2} \partial_T \partial_{\bar{T}} Z_{torus}. \quad (14)$$

Using the relation $\partial_T \partial_{\bar{T}} Z_{torus} = \frac{4\tau_2}{(T+\bar{T})^2} \partial_\tau \partial_{\bar{\tau}} (\tau_2 Z_{torus})$, the integral (14) becomes a boundary integral getting contributions only from massless states. The result is:

$$\partial_{\bar{T}} \Theta_T^{gr} - \partial_T \Theta_{\bar{T}}^{gr} = \frac{1}{16\pi^2} \frac{i\hat{b}^{gr}}{(T+\bar{T})^2}, \quad (15)$$

which finally yields: $\Delta^{gr} = \frac{\hat{b}^{gr}}{32\pi^2} \log(T+\bar{T}) |\eta(iT)|^4$.

In the case of gauge couplings the constant $\hat{b}$ is identified with the β -function of the $N=2$ theory, which, by comparing with the CP-even part, is given by the q^0 term of the expression:

$$\hat{b}_i = \lim_{\tau_2 \rightarrow \infty} Tr Q_h^2 \bar{Q}_i^2 q^{H-1} \bar{q}^{\bar{H}-2}. \quad (16)$$

In (16) $\bar{Q}_i$ is the gauge group generator corresponding to the zero mode part of the Kac-Moody current and Q_h is the helicity operator emerging from the integration of $\langle \psi(z) \psi(0) \rangle^2$ in (9). Q_h^2 can therefore be replaced by $\frac{-d}{i\pi d\tau} \log Z_\psi$, where Z_ψ is the contribution of the two trasverse space-time fermionic coordinates. Using that $Z_\psi = -\frac{\theta_2}{\eta}$ or $\frac{\theta_3 \pm \theta_4}{2\eta}$ for spacetime fermions² or bosons, one finds the familiar one loop formula:

$$\hat{b}_i = \frac{1}{12} Tr_S Q_i^2 - \frac{11}{6} Tr_V Q_i^2 + \frac{1}{3} Tr_F Q_i^2, \quad (17)$$

where S,V and F stand for scalars, vectors and fermions respectively.

In the case of the gravitational coupling Q_i^2 is replaced by the right moving "helicity" operator $\bar{Q}_h^2$, which is given by $\frac{d}{i\pi d\tau} \log \bar{\eta}^2$ as discussed above. In this case one finds:

$$\hat{b}^{gr} = (\frac{1}{12} N_S^L - \frac{11}{6} N_V^L + \frac{1}{3} N_F^L) (\frac{1}{6} N_S^R - \frac{11}{3} N_V^R) + c(n_B - n_F), \quad (18)$$

where $N_{S,V,F}^L$ and $N_{S,V,R}^R$ count the corresponding left and right moving degrees of freedom, and the last term proportional to the difference between the total numbers of bosons and fermions has been added, because in the derivation of the above formula

² The minus sign stands for the spin-statistics.

space-time supersymmetry has been used. Using now the relations $N_S^L N_S^R = N_S$, $N_F^L N_S^R = N_F$ and $N_S^L N_V^R + N_V^L N_S^R = N_V$ with N_S, N_F, N_V the number scalars, two-component fermions and vector bosons respectively, and choosing $c = \frac{1}{120}$ one finds:

$$\hat{b}^{gr} = \frac{1}{45}(N_S + \frac{7}{4}N_F - 13N_V - \frac{113}{2}N_F^L N_V^R + 304N_V^L N_V^R). \quad (19)$$

Eq.(19) reproduces correctly the known trace anomaly coefficients³ for scalars, Majorana fermions and vector bosons. Furthermore the coefficient of $N_F^L N_V^R$ counts for the contribution of a gravitino together with a Majorana fermion and it agrees with the trace anomaly of a spin-3/2, which is $-233/4$ in units where a scalar contributes one. Finally, the coefficient of $N_V^L N_V^R$ counts for the contribution of the graviton, dilaton and two index antisymmetric tensor, and it agrees with the trace anomaly of the spin-2 (212), of a scalar (1), and of a rank-two antisymmetric tensor (91) [7].

In conclusion, we have shown that threshold corrections for gravitational couplings, for the orbifold compactification of the heterotic string, have the same structure and the same origin, i.e. the $N = 2$ sector, as those for gauge couplings. Also, as a technical remark, one can notice that the above calculation provides a relatively simple derivation, if compared to field theory case [8], of the trace anomaly coefficients of the various fields coupled to gravity.

Acknowledgements

I.A. and K.S.N. thank ICTP and Ecole Polytechnique respectively, for hospitality during the completion of this work. One of us (I.A.) thanks S.Ferrara for discussions.

³ These are the coefficients of the square of the Riemann tensor in the expansion of the trace of the energy-momentum tensor.

REFERENCES

- [1] V.S.Kaplunovsky, Nucl.Phys.**B 307** (1988),145.
- [2] L.J.Dixon, V.S.Kaplunovsky and J.Louis, Nucl.Phys.**B 355** (1991),649.
- [3] I.Antoniadis, K.S.Narain and T.R.Taylor, Phys.Lett.**B 267** (1991),37.
- [4] J.-P.Derendinger, S.Ferrara, C.Kounnas and F.Zwirner, *On loop corrections to string effective field theories: field dependent couplings and sigma-model anomalies*, preprint CERN-TH.6004/91(1991).
- [5] I.Antoniadis, E.Gava and K.S.Narain,in preparation.
- [6] B.Zwiebach, Phys.Lett.**B 156** (1985),315.
- [7] S.J.Gates, M.Grisaru, M.Roček, W.Siegel,(1983),*Superspace or One Thousand and One Lessons in Supersymmetry*, (Benjamin/ Cummins).
- [8] S.M.Christensen, M.J.Duff, Nucl.Phys.**B 170**[FSI] (1980),480.

MODULI CORRECTIONS TO GRAVITATIONAL COUPLINGS FROM STRING LOOPS

I. Antoniadis¹, E. Gava^{2,3}, K.S. Narain³

1 Centre de Physique Théorique, Ecole Polytechnique, F-91128 Palaiseau, France

2 Istituto Nazionale di Fisica Nucleare-sez. di Trieste, Italy

3 International Centre for Theoretical Physics, I-34136 Trieste, Italy

Abstract

We study moduli dependent threshold corrections to gravitational couplings in the case of the heterotic string compactified on a symmetric orbifold, for untwisted moduli, extending previous analysis on gauge couplings. Like in the gauge case, the contribution comes entirely from the spacetime $N = 2$ sector. As a byproduct, this calculation provides a simple derivation of the trace anomaly coefficients for the different fields coupled to gravity.

February 1992